

THE MARTINGALE COMPARISON METHOD FOR MARKOV PROCESSES

BENEDIKT KÖPFER,* *University of Freiburg*

LUDGER RÜSCHENDORF,* *University of Freiburg*

Abstract

Comparison results for Markov processes w.r.t. function class induced (integral) stochastic orders have a long history. The most general results so far for this problem have been obtained based on the theory of evolution systems on Banach spaces. In this paper we transfer the martingale comparison method, known for the comparison of semimartingales to Markovian semimartingales, to general Markov processes. The basic step of this martingale approach is the derivation of the supermartingale property of the linking process, giving a link between the processes to be compared. This property is achieved using in an essential way the characterization of Markov processes by the associated martingale problem. As a result the martingale comparison method gives a comparison result for Markov processes under a general alternative but related set of regularity conditions compared to the evolution system approach.

Keywords: Ordering of Markov processes; martingale comparison method; evolution systems

2010 Mathematics Subject Classification: Primary 60E15

Secondary 60J25

1. Evolution systems and comparison of Markov processes

Stochastic ordering and comparison results for Markov processes are a basic problem of probability theory. They have a long history and are motivated by a number of applications in a variety of fields (see [Massey \(1987\)](#), [Cox et al. \(1996\)](#), [Daduna and Szekli \(2006\)](#), [Rüschendorf \(2008\)](#), [Krasin and Melnikov \(2009\)](#), [Rüschendorf and Wolf \(2011\)](#), [Rüschendorf et al. \(2016\)](#), [Criens \(2017\)](#) and [Criens \(2019\)](#)). Various approaches ranging from analytic to coupling methods have been developed

A LGFG grant of the state Baden-Württemberg is gratefully acknowledged

* Postal address: Department of Mathematical Stochastics, University of Freiburg, Freiburg, GER

to this aim sometimes in the context of specific models or specific applications. The most general comparison results so far have been obtained based on the theory of evolution systems on Banach spaces (see [Rüschemdorf et al. \(2016\)](#)).

The transition operators $T_{s,t}$, $s \leq t$, of a Markov process X with values in a metric space S are an evolution system on the space of measurable bounded real-valued functions $\mathcal{L}_b(S)$. Since the transition operators are defined by conditional expectations it is possible to consider them also on function spaces different from $\mathcal{L}_b(S)$. In order to stay in the framework of evolution systems, we consider the transition operators on Banach spaces. We also assume that the Banach spaces are spaces of integrable functions in the sense that they are integrable with respect to all conditional laws. Generally a family of bounded linear operators $(T_{s,t})_{s \leq t}$ from a Banach space \mathbb{B} to \mathbb{B} is called an *evolution system* if for all $0 \leq s \leq t \leq u$ it satisfies

1. $T_{s,s} = \text{id}$,
2. $T_{s,u} = T_{s,t}T_{t,u}$.

An evolution system is called strongly continuous if for all $f \in \mathbb{B}$ the \mathbb{B} -valued function $(s, t) \mapsto T_{s,t}f$ is continuous. If the evolution system is time-homogeneous, i.e. it only depends on the duration $t - s$, then $(T_t)_{t \geq 0}$ defined by $T_t f := T_{0,t}f$ is a semigroup. An evolution system $(T_{s,t})_{s \leq t}$ is called a Feller evolution system if it is strongly continuous and maps $C_0(S)$ into itself. If the evolution system maps $C_b(S)$ into itself, we call it a C_b -Feller evolution system. Further, if the transition operators of a Markov process X are a (C_b) -Feller evolution system, X is called a (C_b) -Feller process.

Right generators of evolution systems $(T_{s,t})$ on a Banach space \mathbb{B} are defined by:

$$A_s^+ f := \lim_{h \downarrow 0} \frac{T_{s,s+h}f - f}{h} \quad \text{for all } s \in \mathbb{R}_+.$$

This operator is defined on its domain $\mathcal{D}(A_s^+)$, i.e. for all $f \in \mathbb{B}$ for which the limit exists in norm. Analogously we define the left generators on the domain $\mathcal{D}(A_s^-)$ by

$$A_s^- f := \lim_{h \downarrow 0} \frac{T_{s-h,s}f - f}{h} \quad \text{for all } s \in \mathbb{R}_+ \setminus \{0\}.$$

If we weaken the limit in the definitions of left and right generators to a pointwise limit, then the corresponding operators are called extended pointwise right and left generators (see [Gulisashvili and van Casteren \(2006\)](#)). The generators of an evolution system on a Banach space \mathbb{B} are linear operators on \mathbb{B} . In general the right generator and left generator do not coincide. In [Böttcher](#)

(2014) an explicit example for a Markov process is given whose right and left generators do not coincide. There also, a condition is given to imply equality for the left and right generators.

Evolution systems arise as solutions of homogeneous evolution problems. Let $(T_{s,t})_{s \leq t \leq T}$ be an evolution system on some Banach space \mathbb{B} . For $0 < t$ we set

$$\mathcal{D}_+(t) := \{f \in \mathbb{B}; s \mapsto T_{s,t}f \text{ is differentiable from the right on } (0, t)\}, \quad (1.1)$$

and

$$\mathcal{D}_+^A(s) := \bigcap_{s \leq t} \mathcal{D}(A_t), \quad \mathcal{D}^A(s, t) := \bigcap_{s \leq u \leq t} \mathcal{D}(A_u).$$

The following theorem restates basic connections of evolution systems to their right generators from Gulisashvili and van Casteren (2006) and states some corresponding representation results.

Theorem 1. *Let $(T_{s,t})_{s \leq t \leq T}$ be an evolution system on a Banach space \mathbb{B} with right generators $(A_t^+)_{t \in [0, T]}$. Then the following assertions hold true:*

- (1) *If $(T_{s,t})_{s \leq t \leq T}$ is strongly continuous, then for fixed t the function $u : s \mapsto T_{s,t}f$ with $f \in \mathcal{D}_+(t)$ is a solution to the following final value problem on $(0, t)$:*

$$\begin{cases} \frac{\partial^+}{\partial s} u(s) &= -A_s^+ u(s), \\ \lim_{s \uparrow t} u(s) &= f. \end{cases} \quad (1.2)$$

- (2) *For $f \in \mathcal{D}_+^{A^+}(s)$, for some fixed $0 < s < T$ and for any $s < t < T$ the forward equation holds:*

$$\frac{\partial^+}{\partial t} T_{s,t}f = T_{s,t}A_t^+ f. \quad (1.3)$$

- (3) *Representation results: Let $(T_{s,t})_{s \leq t \leq T}$ be strongly continuous and $f \in \mathcal{D}_+(t)$. Further, assume that the right derivative $\frac{\partial^+}{\partial u} T_{u,t}f$ is integrable on $[s, t]$. Then the following integral representation of the evolution system holds true:*

$$T_{s,t}f - f = \int_s^t A_u^+ T_{u,t}f du.$$

If $f \in \mathcal{D}^{A^+}(s, t)$ and let the right derivative $\frac{\partial^+}{\partial u} T_{s,u}f$ is integrable on $[s, t]$, then it holds that

$$T_{s,t}f - f = \int_s^t T_{s,u}A_u^+ f du. \quad (1.4)$$

A similar integral representation also holds true for $f \in \mathcal{D}^{A^-}(s, t)$ using the left derivative instead of the right derivative.

Theorem 2. *Let $(T_{s,t})_{s \leq t \leq T}$ be an evolution system on a Banach space \mathbb{B} with left generators $(A_t^-)_{t \in (0, T]}$ and define*

$$\mathcal{D}_-(t) := \{f \in \mathbb{B}; s \mapsto T_{s,t} \text{ is differentiable from the left on } (0, t)\}$$

and

$$\mathfrak{F}_t := \left\{ f \in \mathbb{B}; \lim_{h \downarrow 0} T_{t-h, t} f = f \right\}.$$

Then it holds that:

- (1) For $0 < t < T$ and $f \in \mathcal{D}_-(t) \cap \mathfrak{F}_t$, the function $u : s \mapsto T_{s,t} f$ is a solution to the backward equation on $(0, t)$:

$$\begin{cases} \frac{\partial^-}{\partial s} u(s) &= -A_s^- u(s), \\ \lim_{s \uparrow t} u(s) &= f. \end{cases}$$

- (2) If $(T_{s,t})_{s \leq t \leq T}$ is strongly continuous in s , $f \in \mathcal{D}_-(t)$ and the left derivative $\frac{\partial^-}{\partial s} T_{s,t} f$ is integrable on $[s, t]$, then the following integral representation of the evolution system holds:

$$T_{s,t} f - f = \int_s^t A_u^- T_{u,t} f du.$$

For further extensions and properties of the notion of left (right) generators see [van Casteren \(2011\)](#), [Böttcher \(2014\)](#) and [Köpfer \(2019\)](#).

A basic result in the theory of evolution systems is the following integral representation for solutions to an inhomogeneous evolution problem (see [Rüschemdorf et al. \(2016\)](#)). For fixed $s, t \in \mathbb{R}_+$ let $(A_r)_{s \leq r \leq t}$ be a family of linear operators on a Banach space \mathbb{B} and let $G : [s, t] \rightarrow \mathbb{B}$. A function $u : [s, t] \rightarrow \mathbb{B}$, right differentiable on (s, t) and such that $u(r) \in \mathcal{D}(A_r)$ for all $s \leq r < t$, is called a solution to the *inhomogeneous right evolution problem* with boundary condition $f \in \mathbb{B}$ if

$$\begin{aligned} \frac{\partial^+}{\partial r} u(r) &= -A_r u(r) + G(r) \quad \text{for } s < r \leq t, \\ u(t) &= f. \end{aligned}$$

If additionally u is continuous on $[s, t]$ it is called a classical solution to the inhomogeneous right evolution problem.

On the other hand for $u : [s, t] \rightarrow \mathbb{B}$, left differentiable on (s, t) such that $u \in \mathcal{D}^A(s, t)$, u is a solution to the inhomogeneous left evolution problem with boundary condition $f \in \mathbb{B}$ if

$$\begin{aligned} \frac{\partial^-}{\partial r} u(r) &= -A_r u(r) + G(r) \quad \text{for } s < r \leq t, \\ u(t) &= f. \end{aligned}$$

If u is continuous it is called a classical solution.

The representation result is as follows.

Theorem 3. *Let $(T_{s,t})_{s \leq t}$ be a strongly continuous evolution system on a Banach space \mathbb{B} with right generators $(A_t^+)_{t \geq 0}$. For fix $t \in \mathbb{R}_+$ let $F_t, G : [0, t] \rightarrow \mathbb{B}$ be such that*

- (1) *the function $r \mapsto T_{s,r}G(r)$ is integrable on fixed $[s, t]$,*
- (2) *F_t solves the inhomogeneous right evolution problem for the operators $(A_s^+)_{s \leq t}$,*

$$\frac{\partial^+}{\partial r} F_t(r) = -A_r^+ F_t(r) + G(r).$$

Then the following representation holds

$$F_t(s) = T_{s,t}F_t(t) - \int_s^t T_{s,r}G(r)dr.$$

The same representation result also holds true for the inhomogeneous left evolution problem for left generators of a strongly continuous evolution system (see Köpfer (2019)). The representation result is the basic tool for the general comparison theorem for Markov processes by means of evolution systems in Rüschendorf et al. (2016) stating an ordering result of Markov processes w.r.t. function classes \mathcal{F} . Therefore let X and Y be Markov processes with corresponding transition operators T^X and T^Y . Under some regularity conditions this result states that a propagation of order property for X , i.e. $f \in \mathcal{F}$ implies $T_{s,t}^X f \in \mathcal{F}$ and comparison of generators implies the stochastic ordering condition $X_t \leq_{\mathcal{F}} Y_t$, $t \geq 0$.

The following is a modification of this result holding true for single functions f . Note that for this case where $\mathcal{F} = \{f\}$ the propagation of order property does not make sense. We denote by $\mathcal{D}_+^X(t)$ and $\mathcal{D}_+^Y(t)$ the sets from (1.1) corresponding to the particular transition operators.

Theorem 4. *Assume that $(T_{s,t}^X)_{s \leq t}$ and $(T_{s,t}^Y)_{s \leq t}$ are strongly continuous evolution systems on a Banach space \mathbb{B} and let $f \in \mathcal{D}_+^X(t) \cap \mathcal{D}_+^Y(t)$. If for fixed $t \in \mathbb{R}_+$ it holds that for all $s \leq t$*

- (1) $T_{s,t}^X f \in \mathcal{D}(A_s^{Y+})$;
- (2) $r \mapsto T_{s,r}^Y(A_r^{X+} - A_r^{Y+})T_{s,r}^X f$ is integrable on $[s, t]$;
- (3) $A_s^{X+}T_{s,t}^X f \leq A_s^{Y+}T_{s,t}^X f$ a.s..

Then it holds that

$$T_{s,t}^X f \leq T_{s,t}^Y f \quad \text{a.s. for all } s \leq t.$$

Proof. Set $F_t(s) := T_{s,t}^Y f - T_{s,t}^X f$. Then F_t satisfies by (1.2) the equation

$$\frac{\partial^+}{\partial s} F_t(s) = -A_s^{Y+}T_{s,t}^Y f + A_s^{X+}T_{s,t}^X f,$$

with boundary condition $F_t(t) = 0$. This equation can be written as

$$\frac{\partial^+}{\partial s} F_t(s) = -A_s^{Y+}(T_{s,t}^Y f - T_{s,t}^X f) + (A_s^{X+} - A_s^{Y+})T_{s,t}^X f =: -A_s^{Y+}F_t(s) + G(s), \quad (1.5)$$

where $G(s) := (A_s^{X+} - A_s^{Y+})T_{s,t}^X f$. The terms in the equation are well defined by Theorem 1 and assumption (1). Hence, F_t solves an inhomogeneous right evolution problem.

From the strong continuity of the evolution systems we deduce that F_t is continuous in s . Hence, F_t is a classical solution to the inhomogeneous right evolution problem (1.5).

We show that F_t is nonnegative; then the assertion follows. To see this we apply the integral representation in Theorem 3 to F_t to obtain

$$F_t(s) = T_{s,t}^Y F_t(t) - \int_s^t T_{s,r}^Y G(r) dr = \int_s^t T_{s,r}^Y (-G(r)) dr.$$

From assumption (3) it follows that $-G(r) \geq 0$ a.s. and hence the assertion follows from the fact that the transition operators of Markov processes are positivity preserving operators. \square

A similar comparison result also holds for left generators (see Köpfer (2019)).

2. The martingale comparison method for Markov processes

For the comparison of a semimartingale X to a Markovian semimartingale Y Gushchin and Mordecki (2002) introduced the martingale comparison method. The basic step of this approach is

to establish that the *linking process*

$$(T_{s,t}^X f(Y_s))_{0 \leq s \leq t}$$

is a submartingale for fixed t . Note that $T_{t,t}^X f(Y_t) = f(Y_t)$ and $T_{0,t}^X f(x_0) = E[f(X_t)]$ assuming that $X_0 = x_0 = Y_0$. Thus $(T_{s,t}^X f(Y_s))$ gives a link between the processes X and Y . From the submartingale property of the linking process as a direct consequence the following comparison result is obtained:

$$E[f(Y_t)] = E[T_{t,t}^X f(Y_t)] \geq T_{0,t}^X f(x_0) = E[f(X_t)]. \quad (2.1)$$

If $(T_{s,t}^X f(Y_s))_{0 \leq s \leq t}$ is a supermartingale, the reverse inequality holds. The proof of the submartingale property is essentially based on Itô's formula and on a version of Kolmogorov backwards equation for Markovian semimartingales. In this paper we transfer this martingale comparison approach to the comparison of general Markov processes. As main tool we make essential use of the characterization of Markov processes by the martingale problem. We transfer this classical result (see e.g. [Ethier and Kurtz \(2005, Ch.4., Prop. 1.7\)](#)) to the frame of Markov processes with transition operators defined on a Banach spaces \mathbb{B} of integrable functions; for detailed exposition see [Köpfer \(2019\)](#).

Theorem 5. *Let $(X_t)_{t \in [0, T]}$ be a Markov process with strongly continuous transition operators $(T_{s,t})_{s \leq t \leq T}$ on some Banach space \mathbb{B} and corresponding right generators $(A_t^+)_{t \in [0, T]}$. If for $f \in \mathcal{D}_+^{A^+}(0)$ the right derivative $\frac{\partial^+}{\partial t} T_{s,t} f$ is integrable on $[0, T)$, then it holds that the process $(M_t)_{t \in [0, T]}$ defined by*

$$M_t := f(X_t) - f(X_0) - \int_0^t A_s^+ f(X_s) ds$$

is a martingale with respect to its natural filtration $\mathcal{F}_t := (\sigma(X_s; s \leq t))_{t \in [0, T]}$.

Proof. The integrability is clear since the Banach space is assumed to consist of integrable functions and the right generator $(A_t^+)_{t \in [0, T]}$ maps the Banach space into itself. Let $0 \leq s \leq t$,

then we have by the Markov property and equation (1.4)

$$\begin{aligned}
E[M_t|\mathcal{F}_s] &= E[f(X_t)|X_s] - f(X_0) - \int_0^t E[A_u^+ f(X_u)|X_s] du \\
&= T_{s,t}f(X_s) - f(X_0) - \int_s^t T_{s,u}A_u^+ f(X_s) du - \int_0^s A_u^+ f(X_u) du \\
&= f(X_s) - f(X_0) - \int_0^s A_u^+ f(X_u) du \\
&= M_s.
\end{aligned}$$

This shows the assertion. \square

Remark 1. The integrability condition on $\frac{\partial^+}{\partial t}T_{s,t}f$ can be replaced by the assumption that $T_{s,\cdot}f$ is absolutely continuous on $[0, T)$, see [Hewitt and Stromberg \(1965, Example 18.41\)](#). This also holds true for theorems [4](#), [6](#), [7](#) and [8](#).

A similar martingale property also holds for the left generators. We also will make use of the martingale property for space time functions $f(t, x)$. To that aim we state the following definition, a variant of the definition in [van Casteren \(2011\)](#) for general Banach spaces. The family of operators $(A_t^+)_{t \in [0, T)}$ is here regarded as single operator A^+ on a space consisting of space-time functions $f : [0, T) \times S \rightarrow \mathbb{R}$. Therefore it is important that the Banach space \mathbb{B} on which each A_t^+ is defined can be extended reasonably to functions of the space time process $\bar{\mathbb{B}}$, like $L^p(\mathbb{R}^d)$, $\mathcal{L}_b(\mathbb{R}^d)$ and the smooth functions vanishing at infinity $C_0^\infty(\mathbb{R}^d)$. Then a function f is in $\mathcal{D}(A^+)$ if for all $t \in [0, T]$ it holds that $x \mapsto f(t, x) \in \mathcal{D}(A_t^+)$ and if $(s, x) \mapsto A_s^+ f(s, \cdot)(x) \in \bar{\mathbb{B}}$. We denote by $\mathcal{D}_+(A^+)$ the set of functions $f \in \mathcal{D}(A^+)$ which are right differentiable in the time variable. $\mathcal{D}_-(A^+)$ is defined analogously for the left derivatives.

Definition 1. A family of operators $(A_t^+)_{t \in [0, T)}$ on some Banach space \mathbb{B} is said to be a right generator of a Markov process X if for all $f \in \mathcal{D}_+(A^+)$, for all $x \in S$ and for all $s \leq t$ it holds that

$$\frac{\partial^+}{\partial t} E[f(t, X_t)|X_s = x] = E \left[\frac{\partial^+}{\partial t} f(t, X_t) + A_t^+ f(t, \cdot)(X_t) \middle| X_s = x \right].$$

A family of operators $(A_t^-)_{t \geq 0}$ on \mathbb{B} is said to be a left generator of X if we replace the right derivatives above by left derivatives.

We remark that also the extended pointwise right and left generators (A_t^+) and (A_t^-) of strongly continuous transition operators are right and left generators in the sense of [Definition 1](#) (see [Köpfer \(2019\)](#)).

With the help of this definition we can formulate the martingale property for the space-time process.

Theorem 6. *Let $(A_t^+)_{t \in [0, T]}$ be the right generator of a Markov process X in the sense of Definition 1. Then for every $f \in \mathcal{D}_+(A^+)$ such that $\frac{\partial^+}{\partial t} E[f(t, X_t) | X_s]$ is integrable on $[0, T]$, we have that*

$$M_t := f(t, X_t) - f(0, X_0) - \int_0^t \left(\frac{\partial^+}{\partial s} + A_s^+ \right) f(s, X_s) ds$$

is a martingale with respect to the natural filtration $(\mathcal{F}_t)_{t \in [0, T]}$.

Proof. Again the integrability is clear and the martingale property can be shown straightforward:

$$\begin{aligned} E[M_t | \mathcal{F}_s] &= E[f(t, X_t) | X_s] - f(0, X_0) - \int_0^t E \left[\left(\frac{\partial^+}{\partial u} + A_u^+ \right) f(u, X_u) \middle| X_s \right] du \\ &= E[f(t, X_t) | X_s] - f(0, X_0) - \int_s^t E \left[\left(\frac{\partial^+}{\partial u} + A_u^+ \right) f(u, X_u) \middle| X_s \right] du \\ &\quad - \int_0^s \left(\frac{\partial^+}{\partial u} + A_u^+ \right) f(u, X_u) du \\ &= E[f(t, X_t) | X_s] - f(0, X_0) - \int_s^t \frac{\partial^+}{\partial u} E[f(u, X_u) | X_s] du \\ &\quad - \int_0^s \left(\frac{\partial^+}{\partial u} + A_u^+ \right) f(u, X_u) du \\ &= E[f(t, X_t) | X_s] - f(0, X_0) - E[f(t, X_t) | X_s] + f(s, X_s) \\ &\quad - \int_0^s \left(\frac{\partial^+}{\partial u} + A_u^+ \right) f(u, X_u) du \\ &= M_s. \end{aligned}$$

This completes the proof. □

Remark 2. 1. In [van Casteren \(2011\)](#) a similar result is given under the assumption that the function f is continuously differentiable in the time variable.

2. The proof of Theorem 6 can also be adapted for the extended pointwise right and left generators of the transition operators of X . Thus, Theorem 5 also holds for the extended pointwise right and left generators.

The connection of Markov processes to martingales allows the introduction of further extensions of generators. Therefore, we give some definitions which are motivated by Theorem 5. They are variants of definitions from [Çinlar et al. \(1980\)](#) for time-inhomogeneous Markov processes.

Definition 2. Let $(A_t)_{t \geq 0}$ be a family of operators on a Banach space with domains $(\mathcal{D}(A_t))_{t \geq 0}$. It is called the *extended generator* of a Markov process X if $\mathcal{D}_+^A(0)$ consists of measurable functions $f : S \rightarrow \mathbb{R}$ such that for all $t \geq 0$

$$f(X_t) - f(X_0) - \int_0^t A_s f(X_s) ds$$

is well defined and a local martingale.

Note that it makes no sense to distinguish between left and right generators since here the interpretation as partial semi-differential of the underlying evolution system is not taken. Also there is no restriction to Banach spaces as domains.

The same definition can be given for the space-time process. Recall that the Banach space under consideration has to be extendable to the space-time process.

Definition 3. Let $(A_t^+)_{t \geq 0}$ be a family of operators on a Banach space with domains $(\mathcal{D}(A_t^+))_{t \geq 0}$. It is called the *extended right generator* of the space-time process (id, X) if $\mathcal{D}_+(A^+)$ consists of measurable functions $f : \mathbb{R}_+ \times S \rightarrow \mathbb{R}$ such that for all $t \geq 0$

$$f(t, X_t) - f(0, X_0) - \int_0^t \left(\frac{\partial^+}{\partial s} + A_s^+ \right) f(s, X_s) ds$$

is well-defined and a local martingale.

If the derivatives above are replaced by left derivatives we call the corresponding family of operators extended left generators.

The extended generators can be expanded to other integrals than the Lebesgue integral. This is particularly interesting for example if we consider general Markovian semimartingales with fixed jump times.

Definition 4. Let $F \in \mathcal{V}^+$ be predictable. A family of operators $(A_t)_{t \geq 0}$, $A_t : \mathcal{L}(S) \rightarrow \mathcal{L}(\Omega)$ is called *F-random generator* of a Markov process X , if $\mathcal{D}_+^A(0)$ consists of functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$ for which $(A_t f)_{t \geq 0}$ is an optional process such that $Af \cdot F \in \mathcal{V}$ is predictable and

$$f(X_t) - f(X_0) - \int_0^t A_s f dF_s$$

is well-defined and a local martingale.

Based on the martingale problem we obtain a transfer of the martingale comparison method to the comparison of general Markov processes. In the following theorem we consider the transition

operators $T_{s,t}^X$ for fixed t and $f \in \mathbb{B}$ as a function $T_{\cdot,t}^X f : [0, t] \times S \rightarrow \mathbb{R}$. Hence, we can insert the space-time process and obtain the connection to the martingale problem from Theorem 6. Note that we use generators in the sense of Definition 1. For processes X, Y we denote their (right) generators A^{X+} and A^{Y+} . The following comparison result by the martingale method uses similar conditions as the comparison result Theorem 4 by the evolution approach.

Theorem 7. (Comparison by the martingale comparison method.) *Let $(T_{s,t}^X)_{s \leq t}$ and $(T_{s,t}^Y)_{s \leq t}$ be strongly continuous with right generators $(A_s^{X+})_{s \geq 0}$ and $(A_s^{Y+})_{s \geq 0}$. For $f \in \mathbb{B}$ and fixed $t \in \mathbb{R}_+$ assume that for all $s \leq t$ the following holds*

- (1) $T_{\cdot,t}^X f \in \mathcal{D}_+(A^{X+}) \cap \mathcal{D}_+(A^{Y+})$;
- (2) $\frac{\partial^+}{\partial u} E[T_{u,t}^X f(X_u)|X_s]$ and $\frac{\partial^+}{\partial u} E[T_{u,t}^X f(Y_u)|Y_s]$ are integrable on $[0, t]$;
- (3) $\text{supp}(P^{Y_s}) \subset \text{supp}(P^{X_s})$;
- (4) $A_s^{X+} T_{s,t}^X f \leq A_s^{Y+} T_{s,t}^X f$ a.s.

Then it holds that

$$E[f(Y_t)] \geq E[f(X_t)].$$

Proof. For the proof of the assertion, we establish that $(T_{s,t}^X(Y_s))$ is a submartingale. By construction $(T_{s,t}^X f(X_s))_{s \leq t}$ is a martingale; this follows by the Markov property. For $u \leq s$ we have

$$\begin{aligned} E[T_{s,t}^X f(X_s)|\mathcal{F}_u] &= E[E[f(X_t)|X_s]|X_u] \\ &= E[E[f(X_t)|\mathcal{F}_s]|X_u] \\ &= E[f(X_t)|\mathcal{F}_u] \\ &= T_{u,t}^X f(X_u). \end{aligned}$$

On the other hand by assumption (2) and Theorem 6

$$T_{s,t}^X f(X_s) - T_{0,t}^X f(X_0) - \int_0^s \left(\frac{\partial^+}{\partial u} + A_u^{X+} \right) T_{u,t}^X f(X_u) du$$

is by Theorem 6 a martingale as well. It follows that the integral process

$$\int_0^s \left(\frac{\partial^+}{\partial u} + A_u^{X+} \right) T_{u,t}^X f(X_u) du$$

is also a martingale starting in zero. Since it is an integral with respect to the Lebesgue measure, it is of finite variation and continuous. By [Jacod and Shiryaev \(2003, Corollary I.3.16\)](#) it follows that it is zero $\lambda \times P$ almost surely,

$$\left(\frac{\partial^+}{\partial u} + A_u^{X+}\right) T_{u,t}^X f(X_u) = 0.$$

Hence, for all $x \in \text{supp}(P^{X_u})$ except on a set of time points u of Lebesgue measure zero we obtain

$$\left(\frac{\partial^+}{\partial u} + A_u^{X+}\right) T_{u,t}^X f(x) = 0.$$

By assumption (3), this implies that

$$\left(\frac{\partial^+}{\partial u} + A_u^{X+}\right) T_{u,t}^X f(Y_u) = 0 \tag{2.2}$$

$\lambda \times P$ almost surely. Therefore by assumptions (1), (2) and [Theorem 6](#) applied to $T_{s,t}^X f$ we get that

$$M_s := T_{s,t}^X f(Y_s) - T_{0,t}^X f(Y_0) - \int_0^s \left(\frac{\partial^+}{\partial u} + A_u^{Y+}\right) T_{u,t}^X f(Y_u) du$$

is a martingale. Combining this with [\(2.2\)](#) this implies that

$$T_{s,t}^X f(Y_s) - T_{0,t}^X f(x_0) - \int_0^s (A_u^{Y+} - A_u^{X+}) T_{u,t}^X f(Y_u) du$$

is a martingale. By assumption (4) the integral is non-negative and it follows that $(T_{s,t}^X f(Y_s))_{s \leq t}$ has the representation

$$T_{s,t}^X f(Y_s) = T_{0,t}^X f(x_0) + M_s + \int_0^s (A_u^{Y+} - A_u^{X+}) T_{u,t}^X f(Y_u) du.$$

This is a submartingale since the integral is non-negative. The assertion then follows by inequality [\(2.1\)](#). \square

Remark 3. From the proof of [Theorem 7](#) it follows in a similar way that the inverse inequality, $A_s^{X+} T_{s,t}^X f \geq A_s^{Y+} T_{s,t}^X f$, implies that $(T_{s,t}^X f(Y_s))_{0 \leq s \leq t}$ is a supermartingale and hence the expectations are ordered the other way around $E[f(Y_t)] \leq E[f(X_t)]$.

Since we also have an analog martingale property for left generators, we can transfer [Theorem 7](#) to left generators.

Theorem 8. *Let $(T_{s,t}^X)_{s \leq t}$ and $(T_{s,t}^Y)_{s \leq t}$ be strongly continuous and $f \in \mathbb{B}$. For fixed $t \in \mathbb{R}_+$ assume that for all $s \leq t$ we have*

- (1) $T_{\cdot,t}^X f \in \mathcal{D}_-(A^{Y^-}) \cap \mathcal{D}_-(A^{X^-})$;
- (2) $\frac{\partial^-}{\partial u} E[T_{u,t}^X f(X_u)|X_s]$ and $\frac{\partial^-}{\partial u} E[T_{u,t}^X f(Y_u)|Y_s]$ are integrable on $[0, t]$;
- (3) $\text{supp}(P^{Y_s}) \subset \text{supp}(P^{X_s})$;
- (4) $A_s^{X^-} T_{s,t}^X f \leq A_s^{Y^-} T_{s,t}^X f$ a.s.

Then it holds that

$$E[f(Y_t)] \geq E[f(X_t)].$$

Proof. The proof is similar to the proof of Theorem 7. □

The extended generators and the random generators are defined by a local martingale property. Since the results above rely on the martingale property we are able to obtain similar results for the extended generators and the random generators. The main difference now is that we only have a local martingale property and we are not restricted to Banach spaces.

Let $(A_t^{X^+})_{t \geq 0}$ and $(A_t^{Y^+})_{t \geq 0}$ be the extended right generators for X and Y , see Definition 3. Let $f : S \rightarrow \mathbb{R}$ be a function such that $(T_{s,t}^X f)_{s \geq 0} \in \mathcal{D}_+(A^{X^+})$. By the martingale property of $(T_{s,t}^X f(X_s))_{s \geq 0}$ we get that $\lambda \times P$ almost surely

$$\left(\frac{\partial^+}{\partial s} + A_s^{X^+} \right) T_{s,t}^X f(X_s) = 0. \quad (2.3)$$

We then can undertake the same steps as in the proof of Theorem 7 which yields a local submartingale property for $(T_{s,t}^X f(Y_s))_{s \geq 0}$. So we only need to specify the particular assumptions and make sure that $(T_{s,t}^X f(Y_s))_{s \geq 0}$ is a proper submartingale. Therefore we use of the class (DL) which makes a local (sub-) martingale a proper martingale. A stochastic process X is of class (DL) if all $t \geq 0$, $\{X_\tau; \tau \text{ a stopping time, } \tau \leq t\}$ is uniformly integrable. Note that we use the extended right generator in the next theorem.

Theorem 9. *Let $f : S \rightarrow \mathbb{R}$ be such that for fixed t and all $s \leq t$ it holds*

- (1) $T_{\cdot,t}^X f \in \mathcal{D}_+(A^{X^+}) \cap \mathcal{D}_+(A^{Y^+})$,
- (2) $\text{supp}(P^{Y_s}) \subset \text{supp}(P^{X_s})$,
- (3) $(T_{s,t}^X f(Y_s)^-)_{s \geq 0}$ is of class (DL),

$$(4) \quad A_s^{X+} T_{s,t}^X f \leq A_s^{Y+} T_{s,t}^X f \text{ a.s.}$$

Then

$$E[f(Y_t)] \geq E[f(X_t)].$$

Proof. The proof is similar to the proof of Theorem 7. As mentioned above we have $\lambda \times P$ almost surely

$$\left(\frac{\partial^+}{\partial s} + A_s^{X+} \right) T_{s,t}^X f(X_s) = 0.$$

By assumption (2) we obtain that $\left(\frac{\partial^+}{\partial s} + A_s^{X+} \right) T_{s,t}^X f(Y_s) = 0$ $\lambda \times P$ almost surely as well. On the other hand by the definition of the extended generator it holds that

$$T_{s,t}^X f(Y_s) - T_{0,t}^X f(x_0) - \int_0^s \left(\frac{\partial^+}{\partial u} + A_u^{Y+} \right) T_{u,t}^X f(Y_u) du$$

is a local martingale. Combining with (2.3)

$$T_{s,t}^X f(Y_s) - T_{0,t}^X f(x_0) - \int_0^s (A_u^{Y+} - A_u^{X+}) T_{u,t}^X f(Y_u) du.$$

The integral is non-negative by assumption (4) and hence $(T_{s,t}^X f(Y_s))_{0 \leq s \leq t}$ is a local submartingale. By assumption (3) it is a proper submartingale. The assertion now follows as in Theorem 7. \square

An analog result also holds in the case of extended left generators.

Finally, if we consider random generators instead of extended generators, we get a similar comparison result. Here we have the advantage that there appears no partial derivative in Definition 4. This means that we can proceed more directly.

Theorem 10. *Let $F \in \mathcal{V}^+$ be predictable. Assume that X and Y possess F -random generators $(A_t^X)_{t \geq 0}$ and $(A_t^Y)_{t \geq 0}$. Further, for $f \in \mathcal{D}_+^{A^X}(0) \cap \mathcal{D}_+^{A^Y}(0)$ let*

- (1) $X_0 \sim Y_0$,
- (2) $(f(X_t) - f(Y_t))_{t \geq 0}$ be of class (DL),
- (3) $A_s^X f \geq A_s^Y f$ a.s.

Then we obtain

$$E[f(X_t)] \leq E[f(Y_t)].$$

Proof. Since $(A_t^X)_{t \geq 0}$ and $(A_t^Y)_{t \geq 0}$ are F -random generators we have by Definition 4 that

$$f(X_t) - f(X_0) - \int_0^t A_s^X f dF_s$$

and

$$f(Y_t) - f(Y_0) - \int_0^t A_s^Y f dF_s$$

are local martingales. It follows that

$$\begin{aligned} f(X_t) - f(X_0) - \int_0^t A_s^X f dF_s - f(Y_t) + f(Y_0) + \int_0^t A_s^Y f dF_s \\ = f(X_t) - f(X_0) - f(Y_t) + f(Y_0) + \int_0^t (A_s^Y f - A_s^X f) dF_s \end{aligned}$$

is a local martingale as well. The integral is non-negative and it follows that $(f(X_t) - f(X_0) - f(Y_t) + f(Y_0))_{t \geq 0}$ is a local supermartingale and thus by assumption (2) a supermartingale. In consequence we have that

$$E[f(X_t) - f(Y_t) - f(X_0) + f(Y_0)] \leq 0.$$

The assertion follows by assumption (1). \square

Discussion and comparison of regularity conditions

The martingale comparison method as developed in this paper gives an alternative approach to the comparison of two Markov processes. The regularity conditions of this approach are comparable but different from the conditions assumed for the evolution approach. The main achievements and corresponding assumptions in this paper are the following:

- 1) We extend the evolution approach to the comparison of the expectation of single functions f and thus without using the propagation of order condition.
- 2) The martingale approach allows comparison results also for the case of random generators as occurring for example in the case of Markovian semimartingales with fixed jump times, which do not fit with the conditions of the evolution approach.
- 3) In the case where the evolution approach and the martingale approach both apply, the regularity conditions are similar but (slightly) different. The evolution approach result (Theorem 4) makes use of the integrability condition

(I) $T_{s,u}^Y(A_u^{X+} - A_u^{Y+})T_{s,u}^X f$ is integrable over $u \in [s, t]$,

respectively the related absolute continuity condition, see Remark 1. The martingale approach (Theorem 7) makes use of the condition

(II) $\frac{\partial^+}{\partial u} E[T_{u,t}^X f(X_u)|X_s]$ and $\frac{\partial^+}{\partial u} E[T_{u,t}^X f(Y_u)|Y_s]$ are integrable on $[s, t]$.

Assuming exchangability of differentiation and expectation, condition (II) is equivalent to the integrability of $T_{s,u}^X A_u^{X+} T_{u,t}^X f$ and $T_{s,u}^Y A_u^{X+} T_{u,t}^X f$.

Under the propagation of order condition as in previous literature, for a class of functions \mathcal{F} this amounts to

(II)' $T_{s,u}^X A_u^{X+} f$ and $T_{s,u}^Y A_u^{X+} f$ are integrable for $f \in \mathcal{F}$.

In comparison condition (I) together with additionally postulating integrability of both terms separately results in

(I)' $T_{s,u}^Y A_u^{X+} f$ and $T_{s,u}^X A_u^{Y+} f$ are integrable for $f \in \mathcal{F}$.

Thus roughly both conditions are equivalent under the propagation of order condition.

- 4) In Theorem 10 we give a reasonable and easily verifiable integrability condition not involving directly the generators of the processes. There is no corresponding result available for the evolution approach.

References

- Böttcher B. (2014). Feller evolution systems: Generators and approximation. *Stochastics and Dynamics, Vol. 14*, No. 3.
- Çınlar E., Jacod J., Protter P., Sharpe M.J. (1980). Semimartingales and Markov processes. *Zeitschrift für Wahrscheinlichkeitstheorie, Volume 54*, 161–219.
- van Casteren J.A. (2011). Markov Processes, Feller Semigroups and Evolution Equations. *World Scientific*.
- Cox J.T., Fleischmann K., Greven A. (1996). Comparison of interacting diffusions and an application to their ergodic theory. *Probability Theory and Related Fields, 105*, 513–528.

- Criens D. (2017). Monotone and convex stochastic orders for processes with independent increments. *ArXiv:1606.04993 [math.PR]*.
- Criens D. (2019). Couplings for processes with independent increments. *Statist. Probab. Lett.* *146*, 161–167.
- Daduna H., Szekli R. (2006). Dependence ordering for Markov processes on partially ordered spaces. *Journal of Applied Probability*, *43*, 793–814.
- Ethier S.N., Kurtz T.G. (2005). Markov Processes, Characterization and Convergence. *John Wiley & Sons*.
- Gulisashvili A., van Casteren J.A. (2006). Non-Autonomous Kato Classes and Feynman–Kac Propagators. *World Scientific*.
- Gushchin A.A., Mordecki E. (2002). Bounds on option prices for semimartingale market models. *Proceedings of the Steklov Institute of Mathematics*, *273*, 73–113.
- Hewitt E., Stromberg K. (1965). Real and Abstract Analysis. *Springer*.
- Jacod J., Shiryaev A.N. (2003). Limit Theorems for Stochastic Processes. *Springer*.
- Köpfer B. (2019). Comparison of stochastic processes by Markov projection and functional Itô calculus. *Phd thesis, Albert-Ludwigs-Universität Freiburg*.
- Krasin V.Yu., Melnikov A.V. (2009). On comparison theorem and its application to finance. *In: Optimality and Risk – Modern Trends in Mathematical Finance, Springer*, 171–181.
- Massey W.A. (1987). Stochastic ordering for Markov processes on partially ordered spaces. *Mathematics of Operations Research*, *12*, 350–367.
- Rüschendorf L. (2008). On a comparison result for Markov processes. *Journal of Applied Probability*, *45*, 279–286.
- Rüschendorf L., Wolf V. (2011). Comparison of Markov processes via infinitesimal generators. *Statistics and Decisions*, *28(2)*, 151–175.
- Rüschendorf L., Schnurr A., Wolf V. (2016). Comparison of time-inhomogeneous Markov processes. *Advances in Applied Probability*, *48*, 1015–1044.